

Durham Research Online

Deposited in DRO:

18 July 2014

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Tuxworth, Luke and Baiget, Lise and Phanopoulos, Andreas and Metters, Owen J. and Batsanov, Andrei S. and Fox, Mark A. and Howard, Judith A. K. and Dyer, Philip W. (2012) 'Phosphine-alkene ligand-mediated alkyl-alkyl and alkyl-halide elimination processes from palladium(II).', *Chemical communications.*, 48 (84). pp. 10413-10415.

Further information on publisher's website:

<http://dx.doi.org/10.1039/c2cc35623f>

Publisher's copyright statement:

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

Phosphine-alkene ligand-mediated alkyl-alkyl and alkyl-halide elimination processes from palladium(II)†

Luke Tuxworth,^{a,b} Lise Baiget,^{a,b} Andreas Phanopoulos,^b Owen J. Metters,^b Andrei S. Batsanov,^b Mark A. Fox,^{a,b} Judith A. K. Howard,^b and Philip W. Dyer^{a,b*}

Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x


N-Diphenylphosphino-7-*aza*-benzobicyclo[2.2.1]hept-2-ene (**2**) behaves as a chelating phosphine-alkene ligand for Pd⁰ and Pd^{II}, promoting direct alkyl-alkyl and indirect alkyl-halide reductive elimination due to the stabilisation of the resulting bis(phosphine-alkene)Pd⁰ complex.

Palladium-mediated catalytic transformations are at the heart of contemporary synthetic organic chemistry, particularly for the controlled formation of C–C and C–heteroatom bonds.^{1,2} Manipulation of these reactions necessitates a detailed understanding of the overall reaction trajectory and of the individual steps involved in the catalytic cycle. In particular, establishing factors that control reductive elimination from palladium is a key objective, since it is often through this step that final product formation occurs, *via* either a direct (four-coordinate) or indirect (three-coordinate) transition state.^{3,4} The rate of reductive elimination is intimately linked to the nature of the coupling partners and to the palladium's ancillary ligands, with both steric and electronic factors playing a significant contribution in each case.⁵ Significant effort has been devoted to studying the impact of a wide variety of ubiquitous monodentate phosphine and bidentate diphosphine metal scaffolds, with both their σ -donor character and chelate bite angle having been found to impact directly on elimination rates.^{6,7,8,9}

Recently, efforts have begun to focus on the use of heteroditopic chelate ligands to increase control and further promote elimination. For example, a phosphine bearing a pendant electron deficient alkene has been shown to favour reductive elimination over a competing β -hydride elimination reaction pathway in Negishi cross-couplings.^{10,11} The system's selectivity was attributed to the presence of the strongly π -accepting alkene moiety.

Since it is now well established that weakly *trans* influencing, poorly-donating/electron-accepting phosphines favour reductive elimination from Pd^{II}, we sought to explore the efficacy of a chelating phosphine-alkene (P-alkene) framework combining a weakly basic R₃P moiety with a strongly π -accepting alkene unit.^{12,13,14} To this end, we describe here our use of a strained bicyclic 7-*aza*-norbornene motif to maximise Pd-to-alkene π -retrodonation through relief

of ring strain.




Scheme 1 Conditions: i) Ph₂PCl, NEt₃, CH₂Cl₂, –30 °C to r.t., 12 h, 69%; ii) Se, CDCl₃, r.t., 15 mins., 100%.

The ligand *N*-Ph₂P-7-*aza*-benzobicyclo[2.2.1]hept-2-ene (**2**) was prepared in 69% yield from 7-*aza*-benzobicyclo[2.2.1]hept-2-ene (**1**), *via* a straightforward nucleophilic substitution strategy, Scheme 1.^{15,16,†} Compound **2** presents a single resonance by ³¹P{¹H} NMR spectroscopy, $\delta^{31}\text{P}\{\text{H}\}$ 41.6 ppm, typical of an aminophosphine.¹⁷ The barrier to inversion at nitrogen in compound **2** has been determined computationally (B3LYP/6-31G*) to be low, ~3 kcal mol^{–1} (*cf.* 16 kcal mol^{–1} computed for amine **1**), permitting **2** to adopt the necessary conformation for κ^2 -P,C P-alkene metal chelation. The magnitude of ¹J_{Se-P} (792 Hz) for the phosphine-selenide **2**.Se confirms that the phosphorus donor moiety of **2** is weakly basic.^{17,†}

To probe the ability of P-alkene **2** to enhance reductive elimination from Pd^{II}, its reaction with PdMe₂ and PdCl(Me) fragments has been explored. Treating a toluene solution of PdMe₂(tmeda) with one equivalent of compound **2** cleanly affords *cis*-[PdMe₂(κ^2 -P,C-**2**)] (**3**), which exhibits a single resonance ($\delta^{31}\text{P}\{\text{H}\}$ 81.2 ppm), consistent with P-Pd binding. Coordination of the olefinic moiety is reflected in a shift to lower frequency of the alkene carbon resonance (**2**: δ 144.3 ppm; **3**: δ 119.2 ppm).

Complex **3** is thermally unstable in solution and over 5 days at r.t. smoothly evolves to afford half an equivalent of the monometallic Pd⁰ complex [Pd(κ^2 -P,C-**2**)₂] (**4**) ($\delta^{31}\text{P}\{\text{H}\}$ +82.3 ppm) and ethane ($\delta^1\text{H}$ 0.82 (s) ppm) as the only organic product, according to ¹H and ³¹P NMR spectroscopy, accompanied by the precipitation of elemental palladium (Scheme 2).§ By comparison, complete thermolysis of [PdMe₂(dmpe)] is only achieved upon heating at 90 °C for 1 week,^{18,19} highlighting the potential of ligand **2** in promoting reductive elimination processes.




Scheme 2 Conditions: i) toluene, r.t., >99%; ii) toluene, r.t., 120 h, >99%.

The X-ray molecular structures of complexes **3** and **4** (Fig. 1)† confirm κ^2 -P,C chelation of **2**. In complex **3** the palladium centre is planar, with one site occupied by the midpoint (Md) of the η^2 -bound C(2)=C(3) bond, which is inclined by 88.3° to the coordination plane. The bond distances to palladium fall within the range observed for other complexes of the type $\text{cis-[PdCl}_2(\eta^2\text{-C=C)(PR}_3\text{)]}$.^{20,21,22,23,24} Consistent with moderate $\text{Pd} \rightarrow \pi^*(\text{C=C})$ retro-donation, the C(2)=C(3) bond in **3** (1.359(3) Å) is longer than the corresponding bond in an uncoordinated analogue of **2**, *N*-BOC-*aza*-benzonorbornadiene oxadisilole (1.315(4) Å).²⁵

In complex **4**, the Pd coordination geometry is distorted-tetrahedral, with the Md-Pd-P planes of the two P-alkene ligands forming an 89.1° dihedral angle. Although the η^2 -coordinated C=C bonds of **4** are longer (1.401(2) and 1.406(2) Å) than that of **3**, as would be expected, the difference is only moderate due to the comparatively poor π -donor character of d^{10} palladium species.²⁶

If $\text{PdMe}_2(\text{tmeda})$ is treated instead with **2** in a Pd:2 ratio of 1:2, much faster and quantitative formation of **4** is achieved (5 h, 20 °C) consistent with rapid reductive elimination. Investigation of this reaction by variable temperature NMR spectroscopy (VT NMR) did not reveal any intermediates on the pathway from **3** to **4**. Consequently, we propose that the second equivalent of ligand **2** reacts with complex **3** to form a five-coordinate intermediate **5**, which then undergoes rapid reductive elimination to afford **4** (Scheme 3).



Scheme 3 Conditions: i) toluene, r.t., >99%.

The rate of reductive elimination from complex **5** is likely to be much greater than that from tetracoordinate complex **3** since prior reports have clearly established rate enhancements in reductive elimination with increasing coordination number.²⁷ The intermediacy of a five-coordinate species here is further supported by the observation that treating complex **3** with either 5 mol% of PPh_3 or propene both lead to a significant increase in the rate of reductive elimination of ethane, with formation of **4** being complete within 7 and 30 h, respectively, at 20 °C (*cf.* 120 h in the absence of L-donor ligand).

To probe the role of the alkene moiety of **2** in promoting reductive elimination from **3**, the P-alkane ligand **6** ($\delta^{31}\text{P}\{^1\text{H}\}$ 38.6 ppm), possessing an *aza*-norbornane rather than *aza*-norbornene framework, was prepared from *N*-BOC-7-*aza*-benzobicyclo[2.2.1]heptane.^{28,†} Reaction of two equivalents of **6** with $\text{PdMe}_2(\text{tmeda})$ in toluene forms $\text{cis-[PdMe}_2(\kappa^1\text{-P-6})_2]$ (**7**), $\delta^{31}\text{P}\{^1\text{H}\}$ 61.0 ppm, which cleanly isomerises to the *trans*-diphosphine complex **8** ($\delta^{31}\text{P}\{^1\text{H}\}$ 59.7 ppm,) over a period of 24 h at 20 °C (Scheme 4); no further reaction of **8** is

observed even in the presence of excess **6** at 60 °C. Clearly κ^2 -P,C chelation is vital in retaining the necessary *cis* geometry for reductive elimination from **3**.






Fig. 1 X-ray molecular structures of **3** and **4**; thermal ellipsoids at 50% probability level. Selected bond distances (Å) in **3**: Pd–P 2.2876(6), Pd–C(01) 2.068(3), Pd–C(02) 2.074(3), Pd...C(2) 2.238(2), Pd...C(3) 2.266(2), P–N 1.732(2); in **4**: Pd–P(1) 2.3440(4), Pd–P(2) 2.3341(4), Pd...C(2) 2.1901(14), Pd...C(3) 2.1912(15), Pd...C(12) 2.2095(15), Pd...C(13) 2.1916(14), P(1)–N(1) 1.7470(13), P(2)–N(2) 1.7454(13).



Scheme 4 Conditions: i) toluene, r.t., >99%; ii) toluene, r.t., 24 h, >99%.


Due to significant M–Cl bond polarisation the barrier to alkyl chloride reductive elimination is much greater than that for alkyl-alkyl elimination; consequently, to the best of our knowledge no examples from Pd^{II} have been reported.¹⁴ Therefore we were interested to probe whether P-alkene ligand **2** could also promote alkyl halide elimination. Notably, reaction of **2** with *cis*- $\text{PdCl}(\text{Me})(\text{cod})$ (Pd:2 = 1:1) does indeed give rise to Pd^0 complex **4** (consistent with reductive elimination), which is formed extremely rapidly (reaction complete within seconds at –40 °C according to VT NMR). However, **4** is obtained along with an equimolar quantity of **9** ($\delta^{31}\text{P}\{^1\text{H}\}$ 38.1 ppm), the methyl phosphonium chloride salt of **2**, and leaving unreacted palladium starting material. In contrast, treating *cis*- $\text{PdCl}(\text{Me})(\text{cod})$ with three equivalents of **2** rapidly and quantitatively affords a 1:1 mixture of **4** and **9** as the only products (Scheme 5).



Scheme 5 Conditions: i) toluene, r.t., >99%.

An analogous reaction of *cis*- $\text{PdCl}(\text{Me})(\text{cod})$ with two or three

equivalents of the P-alkane ligand **6** (~5 mins, 20 °C) affords only *trans*-[PdCl(Me)(κ¹-P-**6**)₂] (**10**) (δ ³¹P 6.7 ppm). To account for the formation of **4** and **9** it is proposed that initially [PdCl(Me)(κ²-P,C-**2**)] (**10**) is formed, which undergoes rapid chloride ligand substitution (labilised by the *trans* alkene moiety²⁹) by **2** to form complex **11**. In the presence of free ligand **2**, complex **11** then undergoes *pseudo*-reductive elimination affording phosphonium salt **9** and Pd⁰ complex **4** (Scheme 6). Presumably, methyl-to-phosphorus migration is favoured here by the electron deficient nature of the amido-substituted phosphorus centre. Access to a pathway along which phosphonium salts may be generated is consistent with the formation of Ph₄PI having been observed following Heck coupling reactions involving [PdI(Ph)(PPh₃)₂] precatalysts.³⁰



Scheme 6 Proposed pathway for the formation of phosphonium salt **9**.

In conclusion, by combining a readily prepared, poorly-donating phosphine component with a strongly π-accepting alkene moiety in a single bidentate ligand framework, **2**, it has been possible to efficiently promote reductive elimination from Pd^{II} complexes. The effectiveness of ligand **2** is attributed, in part, to its ability to form a stable *bis*(phosphine-alkene)Pd⁰ complex, **4**, together with its *cis*-bidentate coordination. Notably, the inclusion of an electron poor phosphine moiety can also allow access to less common intramolecular *pseudo*-reductive elimination processes involving the coordinated phosphorus centre, resulting in phosphonium salt formation, in what can be regarded as a rare example of reductive elimination of chloromethane from Pd^{II}. Work is on-going to probe the utility of ligands such as **2** in a variety of catalytic applications.

Durham University and the EU (Marie Curie IE Fellowship to LB) are gratefully acknowledged for financial support. Johnson Matthey is thanked for the loan of palladium salts. Dr A.M. Kenwright and Mr I.H. McKeag are thanked for assistance with NMR spectroscopy. Acquisition of certain mass spectrometric data by the EPSRC National Mass Spectrometry Service at Swansea University is acknowledged.

Notes and references

- ^a Centre for Sustainable Chemical Processes, Department of Chemistry, Durham University, South Road, Durham, UK, DH1 3LE. Fax: +44(0)191 334 2150; Tel: +44(0)191 384 4737; E-mail: p.w.dyer@durham.ac.uk
- ^b Department of Chemistry, Durham University, South Road, Durham, UK, DH1 3LE
- † Electronic supplementary information (ESI) available: experimental details and characterization data, including crystallographic data for **3** and **4**; computational studies. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/abcdefg
- § Integration of both the ¹H and ³¹P{¹H} NMR spectra was quantified using an internal standard.
- ‡ Bruker SMART 6000 CCD area detector, Mo-K_α radiation (λ=0.71073 Å), T=120 K, Olex2 software.³¹ Crystal data: **3**, C₂₄H₂₄NPPd, M= 463.81,

monoclinic, space group P2₁/c (no. 14), a= 10.2185(3), b= 10.6113(4), c= 19.2717(7) Å, β=104.176(14)°, U= 2026.0(1) Å³, Z=4, μ=1.00 mm⁻¹, 26057 reflections (2θ≤60°), R_{int}=0.061, R=0.031 on 4196 data with I≥2σ(I), wR(F²)=0.061 on all 5912 unique data. CCDC 894607. **4**, C₄₄H₃₆N₂P₂Pd, M= 761.09, triclinic, space group P-1 (No. 2), a=11.9327(6), b=12.2787(6), c=13.2937(6) Å, α=70.057(7), β=74.120(7), γ=81.517(7)°, U=1757.92(7) Å³, Z=2, μ=0.65 mm⁻¹, 40908 reflections (2θ≤70°), R_{int}=0.038, R=0.036 on 11656 data with I≥2σ(I), wR(F²)=0.088 on all 14631 unique data. CCDC 894608.

- J. Tsuji, *Palladium Reagents and Catalysts: New Perspectives for the 21st Century*, 2nd Edtn., Wiley, Chichester, 2004.
- J. F. Hartwig, *Acc. Chem. Res.*, 1998, **31**, 852.
- E. Negishi, *Palladium-Catalyzed Reactions Involving Reductive Elimination*, in *Handbook of Organopalladium Chemistry for Organic Synthesis*, E. Negishi, Ed., Wiley, New York, 2002, Vol. 1, pp. 213.
- A. Dedieu, *Chem. Rev.*, 2000, **100**, 543.
- J. M. Brown and N. A. Cooley, *Chem. Rev.*, 1988, **88**, 1031.
- J. F. Hartwig, *Inorg. Chem.*, 2007, **46**, 1936.
- D. A. Redfield, J. H. Nelson, R. A. Henry, D. W. Moore and H. B. Jonnassen, *J. Am. Chem. Soc.*, 1974, **96**, 6298.
- P. Dierkes and P. W. N. M. van Leeuwen, *J. Chem. Soc., Dalton Trans.*, 1999, 1519.
- S. M. Reid, J. T. Mague and M. J. Fink, *J. Am. Chem. Soc.*, 2001, **123**, 4081.
- X. Luo, H. Zhang, H. Duan, Q. Liu, L. Zhu, T. Zhang and A. Lei, *Org. Lett.*, 2007, **9**, 4571.
- H. Zhang, X. Luo, K. Wongkhan, H. Duan, Q. Li, L. Zhu, J. Wang, A. S. Batsanov, J. A. K. Howard, T. B. Marder and A. Lei, *Chem. Eur. J.*, 2009, **15**, 3823.
- E. Negishi, T. Takahashi and K. Akiyoshi, *J. Organomet. Chem.*, 1987, **334**, 181.
- M. Yamashita, J. V. C. Vicario and J. F. Hartwig, *J. Am. Chem. Soc.*, 2003, **125**, 16347.
- P. K. Sajith and C. H. Suresh, *Inorg. Chem.*, 2011, **50**, 8085.
- M. Lautens, K. Fagnou and V. Zunic, *Org. Lett.*, 2002, **4**, 3465.
- A. H. G. P. Prenzel, N. Deppermann and W. Maison, *Org. Lett.*, 2006, **8**, 1681.
- P. W. Dyer, J. Fawcett, M. J. Hanton, R. D. W. Kemmitt, R. Padda and N. Singh, *Dalton Trans.*, 2003, 104.
- W. de Graaf, J. Boersma, W. J. J. Smeets, A. L. Spek and G. van Koten, *Organometallics*, 1989, **8**, 2907.
- J. Krause, G. Cestari, K.-J. Haack, K. Seevogel, W. Storm and K.-R. Pörschke, *J. Am. Chem. Soc.*, 1999, **121**, 9807.
- A. M. Z. Slawin, H. L. Milton, J. Wheatley and J. D. Woollins, *Polyhedron*, 2004, **23**, 3125.
- R. Shintani, W.-L. Duan, T. Nagano, A. Okada and T. Hayashi, *Angew. Chem. Int. Ed.*, 2005, **44**, 4611.
- C. Thoumazet, L. Ricard, H. Grützmacher and P. Le Floch, *Chem. Commun.*, 2005, 1592.
- M. Ma, S. A. Pullarkat, Y. Li and P.-H. Leung, *Inorg. Chem.*, 2007, **46**, 9488.
- L. Bettucci, C. Bianchini, W. Oberhauser, M. Vogt and H. Grützmacher, *Dalton Trans.*, 2010, **39**, 6509.
- Y.-L. Chen, J.-Q. Sun, X. Wei, W.-Y. Wong, and A. W. M. Lee, *J. Org. Chem.*, 2004, **69**, 7190.
- C. A. Tolman, W. C. Seidel and D. H. Gerlach, *J. Am. Chem. Soc.*, 1972, **94**, 2669.
- K. Tatsumi, A. Nakamura, S. Komiya, A. Yamamoto and T. Yamamoto, *J. Am. Chem. Soc.*, 1984, **106**, 8181.
- T. Ohwada, S. Ishikawa, Y. Mine, K. Inami, T. Yanagimoto, F. Karaki, Y. Kabsawa, Y. Otani and M. Mochizuki, *Bioorg. Med. Chem.*, 2011, **19**, 2726.
- T. P. Cheeseman, L. Odell and H. A. Raethel, *Chem. Commun.*, 1968, 1496.
- M. Sakamoto, I. Schimizu, A. Yamamoto, *Chem. Letts.*, 1995, 1101.
- O. V. Dolomanov, L. J. Bourhis, R. J. Gildea, J. A. K. Howard and H. Puschmann, *J. Appl. Cryst.*, 2009, **42**, 339.